

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1. AGENCY USE ONLY (Leave Blank)

2. REPORT DATE

25 June 1999

3. REPORT TYPE AND DATES COVERED
Technical 6/1/98-5/31/99

4. TITLE AND SUBTITLE

Assembly of Long Chain Phosphatidylcholines at a Liquid-Liquid Interface

5. FUNDING NUMBERS

N00014-89-J-1261

6. AUTHOR(S)

B. L. Smiley and G. L. Richmond

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Dept. of Chemistry
University of Oregon
Eugene, OR 97403

8. PERFORMING ORGANIZATION REPORT NUMBER

ONR Technical Report #28

9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)

Dr. Peter Schmidt
Office of Naval Research Physical Science and Technology, ONR 331
800 North Quincy Street
Arlington, VA 22217-5000

10. SPONSORING/MONITORING AGENCY

Office of Naval Research

11. SUPPLEMENTARY NOTES

Biospectroscopy, submitted May 1999

12A. DISTRIBUTION / AVAILABILITY STATEMENT

Approved for public release: distribution unlimited

12B. DISTRIBUTION CODE

13. ABSTRACT (Maximum 200 words)

Please see attached abstract

19990628 088

14. SUBJECT TERMS

Vibrational Sum Frequency Generation (VSFG); oil-water emulsion systems

15. NUMBER OF PAGES

21

16. PRICE CODE

17. SECURITY CLASSIFICATION
OF REPORT

Unclassified

18. SECURITY CLASSIFICATION
OF THIS PAGE

Unclassified

19. SECURITY CLASSIFICATION
OF ABSTRACT

Unclassified

20. LIMITATION OF ABSTRACT

OFFICE OF NAVAL RESEARCH

Grant N00014-89-J-1261

R&T Code 4131038

ONR Technical Report ONR Technical Report #28

Assembly of Long Chain Phosphatidylcholines at a Liquid-Liquid Interface
by

B. L. Smiley and G. L. Richmond

Biospectroscopy, submitted May 1999

Department of Chemistry
1253 University of Oregon
Eugene, OR 97403

June 1999

Reproduction in whole, or in part, is permitted for any purpose of the United States Government.

This document has been approved for public release and sale; its distribution is unlimited

Assembly of Long Chain Phosphatidylcholines at a Liquid-Liquid Interface

B.L. Smiley and G.L. Richmond*
Department of Chemistry
University of Oregon, Eugene, OR 97403
richmond@oregon.uoregon.edu
phone: (541) 346-4635; fax: (541) 346-5859

Abstract

The molecular-level organization of mixed and pure saturated symmetric chain 1,2-diacyl-*sn*-glycero-3-phosphocholines (PCs) adsorbed at a carbon tetrachloride-aqueous interface is explored by probing the hydrocarbon chain conformation within the adsorbed layer. PCs of the chain lengths found most frequently in biological systems, which in pure form are seen to form either very well-ordered or disordered layers, were observed in these studies to assemble into interfacial layers ranging between disordered and ordered states when mixed in various proportions. Independently, while C₁₆ and shorter chain PCs tend to form disordered layers, a strong increase in ordering is observed for C₁₈ and longer chain PCs, in which the hydrocarbon chains are found to be primarily in an all *trans* conformation. Pure C₁₇-PCs adsorbed at the interface produce layers with an intermediate degree of chain ordering. The ability to tune interfacial layer properties in mixed systems as a function of molecular composition, including PC chain length as demonstrated here, is an important mechanism by which surface characteristics of oil-water emulsion systems can be controlled both *in vivo* and in numerous commercial applications.

Assembly of Long Chain Phosphatidylcholines at a Liquid-Liquid Interface

B.L. Smiley and G.L. Richmond*
Department of Chemistry
University of Oregon, Eugene, OR 97403
richmond@oregon.uoregon.edu
phone: (541) 346-4635; fax: (541) 346-5859

Abstract

The molecular-level organization of mixed and pure saturated symmetric chain 1,2-diacyl-*sn*-glycero-3-phosphocholines (PCs) adsorbed at a carbon tetrachloride-aqueous interface is explored by probing the hydrocarbon chain conformation within the adsorbed layer. PCs of the chain lengths found most frequently in biological systems, which in pure form are seen to form either very well-ordered or disordered layers, were observed in these studies to assemble into interfacial layers ranging between disordered and ordered states when mixed in various proportions. Independently, while C₁₆ and shorter chain PCs tend to form disordered layers, a strong increase in ordering is observed for C₁₈ and longer chain PCs, in which the hydrocarbon chains are found to be primarily in an all *trans* conformation. Pure C₁₇-PCs adsorbed at the interface produce layers with an intermediate degree of chain ordering. The ability to tune interfacial layer properties in mixed systems as a function of molecular composition, including PC chain length as demonstrated here, is an important mechanism by which surface characteristics of oil-water emulsion systems can be controlled both *in vivo* and in numerous commercial applications.

1. Introduction

Oil-water emulsions stabilized by interfacially adsorbed surfactants are utilized extensively in commercial processes and preparations. However, these emulsions also provide an available model system of particular interest to biologists since the oil-water interface between natural emulsion particles in the form of dietary fats and surrounding biological fluids plays a crucial role in metabolic processes [1-3]. The oil-water interface and the action of common natural surfactants such as phosphatidylcholines (PCs) within this unique environment have not been well characterized. PCs, commonly referred to as lecithins

when present as a complex mixture of different hydrocarbon chain structures in food preparations, seem to have been characterized somewhat nonselectively in this context. Careful experimental investigations of PCs and other membrane phospholipids at an oil-water interface are sparse [1, 4-9]. PCs as the majority component of membrane bilayers *in vivo* have been well studied, but adsorbed PC monolayers, which, for example, stabilize the surface of plasma lipoproteins in the blood in a manner dependent on the hydrocarbon chain composition of the adsorbed PCs [10], have not typically been characterized particularly at the molecular level of organization. The latter is due, at least in part, to a lack of suitable experimental techniques although some theoretical models have been proposed [11, 12].

In the present investigations, we explore the molecular level organization of saturated symmetric chain PCs adsorbed at the macroscopic interface between immiscible aqueous and carbon tetrachloride (CCl_4) phases utilizing a vibrational spectroscopic technique with an intrinsic specificity for surfaces. Average ordering within the adsorbed layers is deduced as a function of chain length based on the PC hydrocarbon chain conformation. We extend our previous investigations of PCs adsorbed at the liquid-liquid interface to include molecules with still longer hydrocarbon chains up to C_{22} which could not reasonably be prepared by our previously described method [13] but are feasible with the method described herein. The results show a strong increase in relative ordering for the longer chain PCs, including those with an odd number of carbons per chain. The chain ordering of selected mixtures of adsorbed PCs is seen to be composition dependent, indicative of natural structural variances present in the large variety of functional biological membrane assemblies suited to a multitude of roles. The trends observed in these studies are distinctly different from those which might be expected based on complimentary results obtained for more simple, nonbiological surfactants.

2. Theory of Vibrational Sum Frequency Generation from Interfaces

Investigation of molecular ordering at buried interfaces, such as the liquid-liquid interface between CCl_4 and water, provides a particular experimental challenge to surface scientists. In the case of membrane lipids such as PC monolayers, a few thermodynamic characterizations based on surface tension measurements [1, 7-9] and images generated by fluorescence microscopy [14, 15] have

provided only clues to the molecular-level ordering. On the theoretical side, computer modeling techniques have postulated possible molecular conformations [11, 12]. Vibrational sum frequency generation (VSFG), used here to probe the acyl chain conformation of PCs adsorbed at the aqueous-CCl₄ interface, is a surface-specific vibrational spectroscopic technique which has demonstrated molecular specificity and the ability to probe surface conformation.

The interfacial specificity of SFG is derived from the symmetry constraints imposed on a second-order process which in this case is an optical process. Typically, when an electromagnetic field is incident on a material, a polarization, P , is induced within the material with a magnitude proportional to the magnitude of the electric field, E . However, sufficiently intense electric fields may produce higher order effects within the medium such that P is no longer a linear function of E . In sum frequency generation, two intense laser beams at frequencies ω_1 and ω_2 are overlapped within a medium thereby inducing a polarization at the sum of the two frequencies, $\omega_{SF} = \omega_1 + \omega_2$. The magnitude of the electric field generated at the sum frequency, E_{SF} , as represented in Equation 1 is proportional to the product of the two incident electric fields and the second order nonlinear susceptibility, $\tilde{\chi}^{(2)}$, characteristic of the material.

$$E_{SF} \propto P_{SF} = \tilde{\chi}^{(2)} E_{VIS} E_{IR} \quad [1]$$

Using the widely applied electric dipole approximation, SFG is symmetry forbidden in isotropic materials including many bulk liquids but necessarily allowed at interfaces where inversion symmetry is broken. It is this break in symmetry which occurs at surfaces from which the interfacial specificity of SFG is derived. By this same argument the sum frequency response is selective for orientationally ordered adsorbates at an interface even in the presence of unadsorbed, randomly ordered molecules in either bulk phase defining that interface. VSFG has been widely applied at many types of solid, liquid and gas interfaces, including buried interfaces, owing to this intrinsic surface specificity [16-24].

The second order nonlinear susceptibility tensor, $\tilde{\chi}^{(2)}$, which characterizes the interface includes both resonant and nonresonant components. These VSFG measurements are sensitive to the resonant component of the susceptibility, $\tilde{\chi}_R^{(2)}$, which is enhanced when one of the laser beams, in our case a tunable IR source, comes into resonance with allowed molecular vibrational transitions. The magnitude of

the resonant response, given by $\tilde{\chi}_R^{(2)} = N\langle\alpha^{(2)}\rangle$, is determined by the number of contributing oscillators, N , and an orientationally averaged distribution of the molecular hyperpolarizability, $\alpha^{(2)}$, for the particular chemical group. The molecular hyperpolarizability, as defined in Equation 2 [24],

$$\alpha^{(2)} = \sum_n \frac{A_n}{\omega_n - \omega_{IR} - i\Gamma_n} \quad [2]$$

is a sum over all molecular vibrational resonances, n , where A_n is an amplitude term proportional to the product of the Raman and IR transition moments. A_n will be zero if either the Raman or IR transition moments is zero such that the particular molecular vibration must be both Raman and IR active to be sum frequency active and contribute to the measured SF signal [25]. ω_n in Equation 2 is defined as the frequency of the particular molecular vibrational transition, ω_{IR} is the frequency of the tunable IR source and Γ_n is the line width of the transition. When the IR source is tuned such that $\omega_{IR} = \omega_n$, $\alpha^{(2)}$ in the above expression becomes large. An additional several orders of magnitude signal enhancement is obtained in all of our VSFG measurements by use of a total internal reflection geometry in which the generated sum frequency response is emitted from the sample interface at the critical angle [26-29].

3. Optical system description

The pump laser for our VSF measurements is a Q-switched Nd:YAG laser producing 12 nanosecond pulses at 10 Hz. The 1064 nm output of the pump laser is divided into two components, one of which is frequency doubled to produce the 532 nm visible beam directed into the sample cell. The remaining 1064 nm light generates IR light via a LiNbO₃ optical parametric oscillator tunable over the 3.2-3.7 μm range which is focused and directed at the sample interface overlapping the larger 532 nm beam spot at the interface. Both the visible and tunable IR sources are incident on the interface from the more optically dense CCl₄ subphase in order to permit use of a total internal reflection geometry. This arrangement is also beneficial as the aqueous phase is strongly absorbing in the IR. The visible beam is incident at approximately its critical angle during alignment and is adjusted in order to maximize the generated VSF response. The SF signal is measured in reflection as a function of the incident IR

frequency using a photomultiplier tube and gated detection electronics preceeded by appropriate optical filters.

4. Materials and method of sample preparation

Deuterium oxide (D_2O), 99.9%, HPLC grade used as the aqueous phase was obtained from Cambridge Isotope Laboratories, Andover, MA and adjusted to pH 7.0 with a 10 mM phosphate buffer (Mallinckrodt). Carbon tetrachloride (CCl_4), 99.9% HPLC grade was from Sigma-Aldrich. All phosphatidylcholines, stated purity greater than 99%, were obtained in powdered form from Avanti Polar Lipids, Alabaster, AL and used as received. Chloroform ($CHCl_3$), 99+%, HPLC grade containing 0.5-1% ethanol as stabilizer was purchased from Sigma-Aldrich. Deuterated chloroform, 99.8% D was also from Cambridge Isotope Laboratories and obtained fresh. All glassware and the sample cells were cleaned by soaking in Nochromix reagent (Fisher) followed by thorough rinsing with organic-free water from a Nanopure filtration system (Barnstead), dried in an oven and allowed to cool to room temperature prior to use.

Sample cells were prepared with D_2O buffer added on top of the CCl_4 subphase to form a complete overlayer. D_2O was used as the aqueous phase rather than H_2O in order to reduce adsorption of the IR pulse energy at the interface which can cause boiling. The prepared sample cell was then allowed to sit for at least two hours before spreading the PC dissolved in chloroform at a concentration between 1 and 2.5 mg/ml at the liquid-liquid interface by gently expelling small drops of the $CHCl_3$ solution from a syringe tip placed underneath the air-water interface, allowing them to fall by gravity to the liquid-liquid interface. Non-quantitative spreading of a portion of the PC dissolved in the $CHCl_3$ droplets then occurred at the liquid-liquid interface during dissolution of the $CHCl_3$ into the CCl_4 subphase. Typically for the longer chain PCs, two sample spreadings separated by at least an hour for the initial spreading solvent to dissipate from the interface were necessary to produce a close-packed interfacial layer at the liquid-liquid interface. Samples prepared using $CDCl_3$ as a spreading solvent were produced as described for samples spread from $CHCl_3$. PC stock solutions in $CDCl_3$ were placed in cold storage between uses and protected from exposure to light to inhibit degradation of the solvent (in the absence of stabilizer).

Interfacial layers of mixed PCs were prepared by spreading a solution of mixed PCs dissolved in CHCl_3 in the manner described for the pure PC samples.

5. Results and discussion

5.1 Dependence of Ordering on Chain Length

VSF measurements of surfactants which select sum frequency active vibrational modes with components normal to the liquid-liquid interface have previously been utilized to characterize the relative degree of ordering of component hydrocarbon chains [13, 22-24, 30]. This characterization is based on a comparison of the strength of the methyl (CH_3) vs. methylene (CH_2) symmetric stretch (SS) modes. Since symmetry requirements necessitate that a vibrational mode be both IR and Raman allowed for SFG to occur at an interface, the CH_2 -SS of well-ordered, all *trans* hydrocarbon chains within an adsorbed interfacial layer is SF inactive [31, 32] and will not be observed in our measurements. By this argument, the CH_2 -SS peak is small for well-ordered chains whether the chosen polarization combinations are selective for vibrational modes oriented either normal to or parallel to the interface. The CH_2 asymmetric stretch is also expected to be sum frequency inactive for all *trans* hydrocarbon chains in a well-ordered monolayer [31, 32]. In layers with disordered hydrocarbon chains, the symmetry requirements are relaxed and the CH_2 -SS appears in the sum frequency spectrum for modes both parallel and perpendicular to the interface. The trend for the CH_3 -SS mode is opposite that of the CH_2 -SS mode. In monolayers with well-ordered, all *trans* hydrocarbon chains, the CH_3 symmetry axes will be aligned in a narrow distribution near normal to the interface, producing a strong SF active CH_3 -SS mode. In layers with disordered hydrocarbon chains, the distribution of orientations for the CH_3 symmetry axes is broader and, on average, will have an orientation with a much larger component parallel to the interface plane. Both these factors serve to reduce the CH_3 -SS mode SF intensity normal to the interface in a disordered interfacial layer. We are therefore able to quantify the relative degree of chain-ordering in our adsorbed PCs by comparison of spectrally distinct CH_3 -SS mode to CH_2 -SS mode intensities normal to the interface plane within our SF spectra. The reduced number density of adsorbate molecules expected for a disordered

interfacial layer serves to non-selectively reduce the observed SF intensity for all vibrational modes in the spectrum and should not impact the peak ratio.

The possibility of PC multilayer formation at an oil/water interface has been raised where PCs are adsorbed from the organic phase to the liquid-liquid interface [5, 6], and also where adsorption occurs from the aqueous phase involving mechanical perturbation of the surface [13]. We do not assume that a single PC monolayer is present at the liquid-liquid interface in these studies. Whether monolayers or multilayers are formed at the liquid-liquid interface, perhaps dependant on the PC hydrocarbon chain length, only those layers which lack inversion symmetry will contribute to the SF signal intensity measured [33]. SFG investigations of Langmuir-Blodgett multilayers in which successive monolayers on a substrate are oriented head to head and tail to tail suggest that the measured SF signal originates only from the top monolayer [31]. Although it is not certain from which layer within an adsorbed multilayer our SF response may originate if this structure is present, it is clear that distinct differences in the overall layer chain ordering are indicated by our measurements.

In Figure 1, SF data are shown for s-polarized sum frequency, s-polarized visible and p-polarized IR fields (i.e. ssp polarization conditions) giving the measured SF intensity as a function of wavenumber for the incident IR field. The data have been normalized according to the incident IR intensity measured before the sample cell. In Figure 1(a), the spectrum for adsorbed 1,2-dipalmitoyl-*sn*-glycero-3-PC (C₁₆-PC) indicates disordered hydrocarbon chains evidenced by the presence of a strong CH₂-SS mode centered at approximately 2850 cm⁻¹, and a relatively weak CH₃-SS mode centered near 2875 cm⁻¹. A peak centered at approximately 2932 cm⁻¹ in this spectrum which is included only for completeness and is not utilized in our analysis is assigned to a methylene asymmetric stretch mode (CH₂-AS). In comparison, the spectra given in Figures 1(b) and 1(c) for adsorbed 1,2-diheptadecanoyl-*sn*-glycero-3-PC (C₁₇-PC) and 1,2-distearoyl-*sn*-glycero-3-PC (C₁₈-PC) layers, respectively, show a relatively small CH₂-SS mode combined with a relatively strong CH₃-SS mode indicating well-ordered hydrocarbon chains. The shoulder at about 2945 cm⁻¹ to the high energy side of the CH₂-AS is believed to arise from a methyl Fermi resonance (CH₃-FR) enhanced by mode mixing with the strong CH₃-SS in these spectra. A similar pattern wherein a strong increase in chain ordering was observed for the C₁₈-PC compared to C₁₆- and

shorter chain PCs was observed in samples adsorbed from aqueous PC suspensions to the CCl_4 -aqueous interface in earlier studies [13]. These results are supported by published data obtained from electrochemical measurements of ion permeability for PCs of various chain lengths adsorbed at a polarized nitrobenzene/water interface in which symmetric chain PCs with 16 or fewer carbon atoms per chain formed liquid-expanded monolayers while those with 18 or more carbon atoms per chain formed monolayers present in a liquid-condensed state at the liquid-liquid interface [4, 5]. Possibility for formation of PC multilayers at sufficient PC concentrations in the bulk nitrobenzene phase is also noted in this work [5]. Agreement between our results and those obtained at the nitrobenzene/water interface by PC adsorption from nitrobenzene [4, 5] suggest that our spreading method may be complemented by adsorption of dissolved PCs from the CCl_4 phase which are not initially spread at the interface. Addition of PCs by spreading in CHCl_3 at the interface may produce a disproportionately high PC concentration in the vicinity of the liquid-liquid interface which subsequently enhances the rate of PC adsorption. X-ray scattering studies of multilamellar PC vesicles at room temperature have likewise indicated a decrease in the cross sectional area occupied per chain with increasing chain length attributed to tighter chain packing [34]. It should also be noted that CHCl_3 is a known anesthetic, as are alcohols and CCl_4 , although the latter as a nonpolar molecular species is significantly less potent [35]. It has been suggested in work investigating the mechanism of anesthetic action that the presence of adsorbed PCs at an interface enhances the interfacial concentration of the anesthetic relative to the neat interface [36, 37].

In order to more quantitatively gauge ordering of the PC hydrocarbon chains, a ratio of integrated intensities of the CH_3 -SS peak to the CH_2 -SS peak calculated from the measured SSP sum frequency spectra was used. As mentioned above, the ratio CH_3 -SS/ CH_2 -SS will be large for well-ordered hydrocarbon chains, and small for disordered chains. Experiments at the air/water interface on a variety of adsorbed surfactants by Bell, et al. [24] obtained peak ratios between 10 and 0.8 for the most and least densely packed monolayers, respectively, as calculated from their SSP spectra. No phospholipid samples were included in the mentioned work. Well-ordered layers of adsorbed PCs at a liquid-liquid interface have demonstrated somewhat lower maximum peak ratios [13], possibly as a consequence of hydrophobic solvent interaction with the hydrocarbon chains. Our calculated peak ratios for adsorbed

layers of C₁₅-PC through C₂₂-PC (including odd number chains) are shown in Figure 2 as a function of chain length. Consistent with the trend shown in Figure 1, it is clear that a large increase in ordering is observed for C₁₇- and longer chain PCs as compared to the C₁₅ and C₁₆-PCs which possess a similar degree of disorder. The chains of the C₁₈- through C₂₂-PCs are all well-ordered to a similar extent, each exhibiting a very small CH₂-SS mode combined with a relatively large CH₃-SS mode as was shown for C₁₈-PC in Figure 1(c). The CH₂-SS peak in these spectra appears to attain a minimum size which is not further reduced at greater chain lengths. It is also noteworthy that ordering for PCs with an odd number of carbon atoms per acyl chain does not differ from the observed trend for the more physiologically relevant PCs with even numbers of carbons per chain. This similarity has been reported also in thermodynamic data [38]. The increasing number of methylene units relative to methyl groups at greater chain lengths, which in disordered layers might be expected to increase the CH₂-SS mode intensity based simply on an increased number density of oscillators with increasing chain length, does not appear to be a factor in this trend. Rather, there appears to be a minimum thickness for the hydrophobic chain layer at which well-ordered layers can be formed. A similar trend has been noted in previous work for layers of asymmetric and symmetric chain PCs at a liquid-liquid interface prepared by a different method [13].

Nonquantitative spreading of the PC from CHCl₃ in the process of forming the adsorbed PC interfacial layers, as noted in Methods, has potential consequences in the interpretation of our sum frequency data. Along the path of the incident IR beam in the sample cell CCl₄ phase prior to the sample spot, PCs dissolved in trace quantities in the CCl₄ subphase will selectively absorb energy near the CH₂-AS and CH₂-SS regions of the spectrum to a much greater extent than in the CH₃-SS region due to the much greater number of CH₂ groups present, although absorption peaks may be shifted from the SF intensity peaks. Subsequently, if IR absorption in the subphase by dissolved PCs is significant, the incident IR intensity near the CH₂ stretches will be selectively reduced. The measured SF CH₂-SS mode intensity, as represented in Equation 1 by E_{SF}^2 , would then be reduced relative to the CH₃-SS mode intensity, resulting in an inappropriately large CH₃-SS/CH₂-SS ratio. Final bulk PC concentrations in the CCl₄ phase were on the order of 10 μ M in these experiments if we assume that bulk phase depletion resulting from interfacial adsorption is negligible. The latter is not necessarily a valid assumption for spread

or partially spread layers at an interface where the relative proportion of surface-adsorbed molecules is significant. Given that PC solubility in CCl_4 should increase with increasing chain length and our data show an increase in ordering ratio with increasing PC chain length, the possible significance of this phenomenon was experimentally investigated.

Immediately prior to and after the sample cell, the IR intensity incident on a prepared C_{20} -PC sample and transmitted from the sample cell after total internal reflection from the liquid-liquid interface was measured as a function of frequency using a standard power meter. Assuming total reflection of the IR at the interface and equal path lengths into and out of the cell, the IR intensity at the sample spot was estimated to be intermediate between the incident and reflected beam intensities. Only slight dips in IR intensity were evident at approximately 2850 and 2930 cm^{-1} corresponding to absorbances from solution phase vibrational resonances of CH_2 symmetric and asymmetric stretches, respectively. A more gradual decrease in reflected IR intensity was observed below 2880 cm^{-1} , and a pronounced decrease in IR absorption was observed above 2975 cm^{-1} . IR absorbance measurements of a CCl_4 sample taken from the cell subphase following the experiment identified the decreasing transmission below 2880 cm^{-1} as resulting from an absorption tail of D_2O monomers dissolved in the CCl_4 . From the same measurement, the pronounced decrease in transmission above 2975 cm^{-1} can be attributed to an absorption tail from the CH vibrational resonance of the CHCl_3 spreading solvent, now dissolved in the CCl_4 . Very small absorption peaks corresponding to CH_2 symmetric and asymmetric stretch vibrational resonances were observed overlapping the D_2O absorption tail, and appeared to be insignificant in comparison. These measurements indicate that specific subphase absorption of the tunable IR energy in the vicinity of the CH_2 -SS is not responsible for the small SF peak intensity observed for this mode in the longer-chain PCs. Rather, the hydrocarbon chains of these PCs appear, in fact, to be well-ordered.

An additional consideration which may impact the ordering of the various adsorbed PCs is the presence of ethanol (as a manufacturer-added stabilizer) in the CHCl_3 spreading solvent. Ethanol has been shown to alter/increase the monolayer/bilayer structural properties, even in trace quantities, and may induce phase separations in mixed systems [39]. In anesthetic dosages, alcohols have also been shown to perturb hydration of the PC headgroup [40], suggesting alterations in membrane structural properties.

Since commercially available CHCl_3 is not available without stabilizer, we prepared spreading solutions of PCs dissolved in deuterated chloroform (CDCl_3), which is available without stabilizers and must be handled accordingly, for comparison with the CHCl_3 results. The chain ordering of PCs spread from CDCl_3 did not appear to differ significantly from those prepared with CHCl_3 . The presence of ethanol in the spreading solvent therefore does not appear to be a factor in the observed ordering.

5.2 Mixed C_{16} & C_{18} PCs

If we now consider only physiologically relevant PCs with an even number of carbon atoms per chain, the position at which the transition from a disordered layer to an ordered layer occurs is between C_{16} -PC and C_{18} -PC, as was shown in Figure 2. In biological membranes, the PC alkyl chains are primarily composed of C_{16} and C_{18} chain lengths, C_{20} chains being the other significant portion [41]. If the length of the hydrocarbon chains can be related to structural stability of the interfacial layer, mixtures of "unstable" C_{16} and "stable" C_{18} -PC chains could conceivably produce an intermediate degree of stability at the liquid-liquid interface. This intermediate degree of ordering between rigid inflexibility and random disorder is reflective of the pseudo-stability and complexity inherent in many biological systems. Although our SFG measurements were done *in situ* under ambient conditions (i.e. room temperature), it has been suggested in previous work that laser heating of the sample spot during SFG measurements under the specified experimental conditions may raise the local interface temperature to between 41 and 44°C [13]. While ordering of the interfacial layer is temperature-dependent, the region of the adsorbed layer being sampled may actually be close to body temperature (37°C) and our results therefore applicable to *in vivo* comparisons.

Results from different mole percent mixtures of C_{16} -PC with C_{18} -PC at the liquid-liquid interface are summarized in Figure 3, where we are again using the $\text{CH}_3\text{-SS}/\text{CH}_2\text{-SS}$ peak ratio calculated from SFG ssp spectra as a quantitative measure of hydrocarbon chain ordering. We assume that either the different chain length PCs mix close to homogeneously in the adsorbed layer or that sampling over time during SFG measurements is representative of the average chain ordering present within the layer. DSC measurements of C_{14} - and C_{16} -PC mixtures incorporated into multilamellar vesicles have indicated that

mixing is not ideal and is less homogeneous with a larger mole fraction of the C₁₆-PC [42]. The calculated peak ratio obtained from VSFG measurements is shown vs. mole percent of C₁₈-PC in Figure 3. The data indicate that up to approximately 30% spread C₁₈-PC, the PC chains are as disordered as in a pure C₁₆-PC sample. At larger percent C₁₈-PC compositions, chain ordering steadily increases until a well-ordered layer is obtained at greater than 80% spread C₁₈-PC. It is noteworthy that at spread 50% C₁₈-PC/50% C₁₆-PC, the hydrocarbon chains are still relatively disordered, rather than at the midpoint between ratios calculated for the two pure monolayers. In this 1:1 mixture, the two methylene unit mismatch between neighboring C₁₆- and C₁₈-PC chains may still prohibit close chain packing. There are several other possible explanations for this, the most obvious being that our measure of the hydrocarbon chain ordering, the ratio of integrated intensities CH₃-SS/CH₂-SS, is a relative measure only and may not be a linear function of the chain ordering. The chain ordering also need not vary in a simple "linear" fashion as a function of composition. Experimentally, another factor which may play into the trend observed in Figure 3 is that the PCs when spread independently from CHCl₃ at the liquid-liquid interface appear to vary in their efficiency of spreading such that the final adsorbed layer composition at the interface may differ proportionately from the composition of the spreading solution. Nonideal mixing within the interfacial layer may also be a factor in some way. Other studies have noted that the rate of formation of a close-packed interfacial layer is slower with increasing chain length [5], consistent with our observations. For PCs with hydrocarbon chain lengths greater than 19 carbon atoms, excess PCs from the second spreading were observed to produce flat, elliptical islands on top of the interfacial layer apparently resulting from an inability of PCs in the CHCl₃ droplets to penetrate the interface and thereby dissolve in the CCl₄ subphase. None of the PCs studied are soluble as monomers in the aqueous phase to a significant extent.

VSFG measurements incorporating chain-deuterated PCs were performed in an attempt to better elucidate the ordering of the particular PCs in the binary mixture and possibly glean information about the relative population of the two PCs adsorbed at the interface from the measured SF intensities. Experiments with mixed C₁₆- and C₁₈-PCs in which one or the other species was deuterated seemed to indicate in complimentary mixtures of d-C₁₈-PC/C₁₆-PC and C₁₈-PC/d-C₁₆-PC of the same molar ratio that the chain ordering, as deduced from the peak area ratio of VSFG SSP spectra, was smaller for both

samples than for the peak ratio obtained in similar samples with only hydrogenated chains. Unfortunately, the signal intensities obtained for either one or the other hydrogenated species in the mixtures was too small relative to the noise level to provide for a reliable quantitative comparison. The small size of the generated signal results from the proportionality of the sum frequency intensity to the square of the number density, N , of oscillators as indicated in the Theory section. The smaller signal intensity is combined with a significantly greater noise level as compared to purely hydrogenated samples possibly as the result of interfacial density fluctuations of the hydrogenated sample component within the sampled area. Reduced intermolecular interferences between oppositely oriented CH_2 groups mixed with CD_2 groups in adjacent hydrocarbon chains is a possible explanation for this observation if the smaller $\text{CH}_3\text{-SS}/\text{CH}_2\text{-SS}$ ratios observed are meaningful. To test this possibility, we investigated 50/50 mixtures of d- $\text{C}_{18}\text{-PC}/\text{C}_{18}\text{-PC}$ and d- $\text{C}_{16}\text{-PC}/\text{C}_{16}\text{-PC}$ adsorbed at the interface. The C_{18} mixtures would be expected to give a relatively small $\text{CH}_2\text{-SS}$ intensity regardless of intermolecular interferences and gave ratios within the error bar for results obtained from the purely hydrogenated $\text{C}_{18}\text{-PCs}$. The SF signal intensities measured from the $\text{C}_{16}\text{-PC}$ mixtures were very small and noisy. These results unfortunately did not provide further insight into the adsorbed binary mixtures of C_{16} - and $\text{C}_{18}\text{-PCs}$. It is also not clear whether intermolecular interferences between oppositely oriented CH_2 groups within the sample layers significantly impact these VSFG measurements of hydrocarbon chain ordering.

6. Conclusions

Distinct differences in interfacial layer ordering of adsorbed symmetric chain PCs have been observed as a function of chain length at an aqueous- CCl_4 interface. C_{18} and longer chain PCs were seen to form extremely well ordered interfacial layers with chains in a predominantly all *trans* conformation, while C_{16} and $\text{C}_{15}\text{-PCs}$ formed layers with disordered chains. The $\text{C}_{17}\text{-PCs}$ produced layers with an intermediate degree of order. These results are in contrast to those reported for interfacially adsorbed alkyl sulfonate monolayers at a similar interface in which increasing chain disorder was noted as a function of increasing length of the component hydrocarbon chains as attributed to a greater number of *gauche* conformers [43]. This indicates that the behavior of PCs adsorbed at a liquid-liquid interface is not

determined mainly by chain solvation forces. We might reasonably expect such differences in molecular packing arrangements between relatively simple charged surfactants displaying interheadgroup repulsion and the structurally more complicated zwitterionic PCs. Mixtures of C₁₆ and C₁₈-PCs adsorbed at the liquid-liquid interface exhibited a systematic increase in chain ordering as a function of C₁₈-PC mole fraction representative of compositional alterations in membrane structure which are needed biologically to control membrane function.

Acknowledgements

The authors thank Dr. R.A. Walker for helpful discussions, and The National Science Foundation (CHE 9531402) and the Office of Naval Research for financial support.

References

- [1] T. Handa, H. Saito and K. Miyajima, Phospholipid monolayers at the triolein-saline interface: Production of microemulsion particles and conversion of monolayers to bilayers, *Biochemistry* 29 (1990) 2884-2890.
- [2] K.W. Miller and D.M. Small, Structure of triglyceride-rich lipoproteins: an analysis of core and surface phases, in A.M.J. Gotto (Ed.), *Plasma Lipoproteins*, Elsevier, New York, 1987, pp. 1-75.
- [3] V. Schumaker and A. Lemberas, Lipoprotein metabolism: Chylomicrons, very-low-density lipoproteins and low density lipoproteins, in A.J. Lusis, J.I. Rotter, and R.S. Sparkes (Eds.) *Molecular Genetics of Coronary Artery Disease. Candidate Genes and Processes in Atherosclerosis*, Karger, New York, 1992, pp. 98-139.
- [4] T. Kakiuchi, M. Kotani, J. Noguchi, M. Nakanishi and M. Senda, Phase transition and ion permeability of phosphatidylcholine monolayers at the polarized oil/water interface, *J. Colloid Interface Sci.* 149 (1992) 279-289.
- [5] T. Kakiuchi, Phospholipid Monolayers and Interfacial Enzyme Reactions of Phospholipases at ITIES, in A.G. Volkov and D.W. Deamer (Eds.) *Liquid-Liquid Interfaces: Theory and Methods*, CRC Press, Boca Raton, 1996, pp. 317-331.
- [6] Y.A. Shchipunov and A.F. Kolpakov, Phospholipids at the oil/water interface: Adsorption and interfacial phenomena in an electric field, *Adv. Colloid Interface Sci.* 35 (1991) 31-138.
- [7] J. Mingins, D. Stigter and K.A. Dill, Phospholipid interactions in model membrane systems. I. Experiments on monolayers, *Biophys. J.* 61 (1992) 1603-1615.

- [8] K. Ogino and M. Onishi, Interfacial action of natural surfactants in oil/water systems, *J. Colloid Interface Sci.* 83 (1981) 18-25.
- [9] J. Mingins, J.A.G. Taylor, B.A. Pethica, C.M. Jackson and B.Y.T. Yue, Phospholipid monolayers at non-polar oil/water interfaces. Part 3- Effect of chain length on phase transitions in saturated di-acyl lecithins at the n-heptane/aqueous sodium chloride interface, *J. Am. Chem. Soc., Faraday Trans. 1* 78 (1982) 323-339.
- [10] T.G. Redgrave, V. Rakic, B.-C. Mortimer and J.C.L. Mamo, Effects of sphingomyelin and phosphatidylcholine acyl chains on the clearance of triacylglycerol-rich lipoproteins from plasma. Studies with lipid emulsions in rats, *Biochim. Biophys. Acta* 1126 (1992) 65-72.
- [11] K.A. Dill and D. Stigter, Lateral interactions among phosphatidylcholine and phosphatidylethanolamine head groups in phospholipid monolayers and bilayers, *Biochemistry* 27 (1988) 3446-3453.
- [12] D. Stigter and K.A. Dill, Lateral interactions among phospholipid headgroups at the heptane/water interface, *Langmuir* 4 (1988) 200-209.
- [13] B.L. Smiley and G.L. Richmond, Alkyl chain ordering of asymmetric phosphatidylcholines adsorbed at a liquid-liquid interface, *J. Phys. Chem.*, submitted.
- [14] M. Thoma and H. Möhwald, Phospholipid monolayers at hydrocarbon/water interfaces, *J. Colloid Interface Sci.* 162 (1994) 340-349.
- [15] H. Möhwald, A. Dietrich, C. Böhm, G. Brezesinski and M. Thoma, Domain formation in monolayers, *Mol. Membrane Biol.* 12 (1995) 29-38.

- [16] H. Ishida, K. Iwatsu, J. Kubota, A. Wada, K. Domen and C. Hirose, In situ SFG spectroscopy of film growth. II. Deposition of formic acid on Ni(110) surface, *J. Chem. Phys.* 108 (1998) 5957-5964.
- [17] C.D. Bain, Sum frequency vibrational spectroscopy of the solid/liquid interface, *J. Chem. Soc., Faraday Trans.* 91 (1995) 1281-1296.
- [18] P. Guyot-Sionnest, J.H. Hunt and Y.R. Shen, Sum-frequency vibrational spectroscopy of a Langmuir film: Study of molecular orientation of a two-dimensional system, *Phys. Rev. Lett.* 59 (1987) 1597-1600.
- [19] C. Hirose, N. Akamatsu and K. Domen, Formulas for the analysis of surface sum frequency generation spectrum by CH stretching modes of methyl and methylene groups, *J. Chem. Phys.* 96 (1992) 997-1004.
- [20] J.H. Hunt, P. Guyot-Sionnest and Y.R. Shen, Observation of C-H stretch vibrations of monolayers of molecules: Optical sum frequency generation, *Chem. Phys. Lett.* 133 (1987) 189-192.
- [21] D. Zhang, J. Gutow and K.B. Eisenthal, Vibrational spectra, orientations, and phase transitions in long chain amphiphiles at the air/water interface: Probing the head and tail groups by sum frequency generation, *J. Phys. Chem.* 98 (1994) 13729-13734.
- [22] J.C. Conboy, M.C. Messmer and G.L. Richmond, Dependence of alkyl chain conformation of simple ionic surfactants on head group functionality as studied by vibrational sum-frequency spectroscopy, *J. Phys. Chem. B* 101 (1997) 6724-6733.

- [23] R.A. Walker, J.C. Conboy and G.L. Richmond, Molecular structure and ordering of phospholipids at a liquid-liquid interface, *Langmuir* 13 (1997) 3070-3073.
- [24] G.R. Bell, C.D. Bain and R.N. Ward, Sum-frequency vibrational spectroscopy of soluble surfactants at the air/water interface, *J. Chem. Soc., Faraday Trans. 92* (1996) 515-523.
- [25] Q. Du, R. Superfine, E. Freysz and Y.R. Shen, Vibrational spectroscopy of water at the vapor/water interface, *Phys. Rev. Lett.* 70 (1993) 2313-2316.
- [26] N. Bloembergen and C.H. Lee, Total reflection in second harmonic generation, *Phys. Rev. Lett.* 19 (1967) 835-7.
- [27] N. Bloembergen, H.J. Simon and C.H. Lee, Total reflection phenomena in second-harmonic generation of light, *Phys. Rev.* 181 (1969) 1261-1271.
- [28] J. Löbau and K. Wolfrum, Sum-frequency spectroscopy in total internal reflection geometry: Signal enhancement and access to molecular properties, *J. Opt. Soc. Am. B* 14 (1997) 2505-2512.
- [29] J.C. Conboy, M.C. Messmer and G.L. Richmond, Investigation of surfactant conformation and order at the liquid-liquid interface by total internal reflection sum-frequency vibrational spectroscopy, *J. Phys. Chem.* 100 (1996) 7617-7622.
- [30] R.A. Walker, J.A. Gruetzmacher and G.L. Richmond, Phosphocholine monolayer structure at a liquid-liquid interface, *J. Am. Chem. Soc.* 120 (1998) 6991-7003.
- [31] N. Akamatsu, K. Domen and C. Hirose, SFG study of two-dimensional orientation of surface methyl groups on cadmium arachidate Langmuir-Blodgett films, *J. Phys. Chem.* 97 (1993) 10070-10075.

- [32] K. Wolfrum and A. Laubereau, Vibrational sum-frequency spectroscopy of an adsorbed monolayer of hexadecanol on water. Destructive interference of adjacent lines, *Chem. Phys. Lett.* 228 (1994) 83-88.
- [33] B.L. Smiley, R.A. Walker, D.E. Gragson, T.E. Hannon and G.L. Richmond, Vibrational sum frequency spectroscopy of surfactants and phospholipid monolayers at liquid-liquid interfaces, in *Laser Techniques for Condensed Phase and Biological Systems*, SPIE Vol. 3273 (1998) 134-144.
- [34] W.-J. Sun, S. Tristram-Nagle, R.M. Suter and J.F. Nagle, Structure of gel phase saturated lecithin bilayers: Temperature and chain length dependence, *Biophys. J.* 71 (1996) 885-891.
- [35] Y.-S. Tsai, S.-M. Ma, H. Kamaya and I. Ueda, Fourier transform infrared studies on phospholipid hydration: Phosphate-oriented hydrogen bonding and its attenuation by volatile anesthetics, *Mol. Pharmacol.* 31 (1987) 623-630.
- [36] Y. Suezaki, A. Shibata, H. Kamaya and I. Ueda, Atypical Langmuir adsorption of inhalation anesthetics on phospholipid monolayer at various compressional states: difference between alkane-type and ether-type anesthetics, *Biochim. Biophys. Acta* 817 (1985) 139-146.
- [37] H.-C. Lin, I. Ueda, S.H. Lin, D.D. Shieh, H. Kamaya and H. Eyring, Surface activities of tertiary amine local anesthetics at air/water interface in the presence and absence of phospholipid monolayers, *Biochim. Biophys. Acta* 598 (1980) 51-65.
- [38] L. Finegold and M.A. Singer, The metastability of saturated phosphatidylcholines depends on the acyl chain length, *Biochim. Biophys. Acta* 855 (1986) 417-420.

- [39] E.S. Rowe, Induction of lateral phase separations in binary lipid mixtures by alcohol, *Biochemistry* 26 (1987) 46-51.
- [40] J.-S. Chiou, C.-C. Kuo, S.H. Lin, H. Kamaya and I. Ueda, Interfacial dehydration by alcohols: Hydrogen bonding of alcohols to phospholipids, *Alcohol* 8 (1991) 143-150.
- [41] P. Yeagle, editor, *The Structure of Biological Membranes*, 1992, CRC Press: Boca Raton.
- [42] P. Garidel and A. Blume, Miscibility of phospholipids with identical headgroups and acyl chain lengths differing by two methylene units: Effects of headgroup structure and headgroup charge, *Biochim. Biophys. Acta* 1371 (1998) 83-95.
- [43] J.C. Conboy, M.C. Messmer and G.L. Richmond, Effect of alkyl chain length on the conformation and order of simple ionic surfactants adsorbed at the D_2O/CCl_4 interface as studied by vibrational sum-frequency spectroscopy, *Langmuir*, in press.

Figure 1: Measured SF intensity as a function of the incident IR wavenumber. Spectra are shown for (a) C₁₆-PC, (b) C₁₇-PC and (c) C₁₈-PC adsorbed at the aqueous-CCl₄ interface for SSP polarization conditions. Lines are provided as a guide to the eye.

Figure 2: Calculated CH₃-SS/CH₂-SS peak ratios vs. PC chain length. Error bars are given based on the variance between multiple samples.

Figure 3: CH₃-SS/CH₂-SS peak ratios calculated for various mole percent mixtures of C₁₈-PC with C₁₆-PC spread at the aqueous-CCl₄ interface. SF intensities were measured under SSP polarization conditions.